



Review Article

# Viruses and viroids in tomato (*Solanum lycopersicum*) and plant growth promoting rhizobacteria as a management alternative

**Erika Janet Zamora-Macorra**, Departamento de Preparatoria Agrícola, \***Norma Ávila-Alistac**, **Erika Lagunes-Fortiz**, Departamento de Fitotecnia. Universidad Autónoma Chapingo, Km 38.5, carretera México-Texcoco, Chapingo, Estado de México. C.P. 56230; **Sergio de los Santos-Villalobos**, Instituto Tecnológico de Sonora, 5 de Febrero 818 Sur, Colonia Centro, Ciudad Obregón, Sonora. C.P. 85000.

**\*Corresponding Author:**

Norma Ávila-Alistac  
alixtac@gmail.com

**Section:**  
Special Number

**Received:**  
30 August, 2023  
**Accepted:**  
12 December, 2023  
**Published:**  
28 December, 2023

**Citation:**  
Zamora-Macorra EJ, Ávila-Alistac N, Lagunes-Fortiz E and de los Santos-Villalobos S. 2023. Viruses and viroids in tomato (*Solanum lycopersicum*) and plant growth promoting rhizobacteria as a management alternative. Mexican Journal of Phytopathology 41(4): 6.  
DOI: <https://doi.org/10.18781/R.MEX.FIT.2023-7>



## ABSTRACT


Viruses and viroids cause several diseases in tomato (*Solanum lycopersicum*) worldwide, generating important economic losses. About 312 viruses and seven viroids have been associated, of which more than 28 are present in Mexico. Therefore, the use of Plant Growth-Promoting Rhizobacteria (PGPR) can be an effective alternative for the management of viruses and viroids. The genera *Pseudomonas*, *Bacillus*, *Azospirillum*, *Anabena* and *Stenotrophomonas* have been implemented against main viruses reported in tomato: *Cucumber mosaic virus* (CMV), *Tobacco mosaic virus* (TMV), *Tomato chlorotic spot virus* (TCSV), *Tomato mottle virus* (ToMoV), *Tomato spotted wilt virus* (TSWV), *Tomato yellow leaf curl virus* (TYLCV), *Potato virus Y* (PVY), *Groundnut bud necrosis virus* (GBNV), with benefits in decreased incidence and severity up to 80 % and yield increase over 40 %. In Mexico, only *Bacillus* has been used. The use of PGPR is a strategy that could mitigate the impact of viral and viroid diseases and can be integrated into integrated management.

**Keywords:** ISR, *Solanum lycopersicum*, PGPR, *Pseudomonas*, *Bacillus*, viruses.

The tomato (*Solanum lycopersicum*), native to South America and currently distributed worldwide, is adapted to tropical and temperate conditions (Hanssen and Lapidot, 2012). It is one of the most cultivated vegetables in both greenhouse and open-field settings (Sánchez-del Castillo *et al.*, 2009). Additionally, it stands as one of the most lucrative and globally consumed crops. In 2021, the global harvest reached 256,770,677 tons (FAO, 2023). In Mexico, the cultivation of tomatoes

holds significant social and economic importance due to foreign exchange earnings and job generation (Hernández-Martínez *et al.*, 2004). However, pathogens and insect pests can compromise its yield and quality (Savary *et al.*, 2019). For instance, diseases caused by viruses and viroids result in economic losses worldwide (Ling and Zhang, 2009; Antignus *et al.*, 2002). In Mexico, there are only records of losses of up to 80% of viral infections comparing the use of agricon with the control (without management) (Ramírez-Rojas, 2006), so it is important to have accurate information on the economic impact of these pathogens, since detections are constant (Figure 1) and are a latent risk for national production. A global economic impact exceeding \$30 billion annually has been estimated for viral diseases affecting economically significant crops (Sastry, 2013a).

Both viruses and viroids can lead to losses in crop production and quality. Therefore, it is crucial to explore management alternatives that ensure direct, effective, and ecologically safe control (Rojas *et al.*, 2018). For decades, chemical pesticides have been employed globally in agriculture to control insect pests and



**Figure 1.** States of the Mexican Republic where viruses and viroids were detected in tomato (See Table 1).

phytopathogens, often applied excessively to both crops and soil, resulting in resistance issues (Karthika *et al.*, 2020). Specifically, in the control of diseases caused by viruses and viroids, the use of agrochemicals has been limited to the control of vector insects. However, in recent years, management strategies have shifted towards seeking environmentally friendly alternatives compatible with the surrounding flora and fauna.

Therefore, a comprehensive understanding of these infectious agents and their interaction with hosts is essential to gain knowledge and propose effective management alternatives. While research has been conducted in recent years on the role of Plant Growth-Promoting Rhizobacteria (PGPR) in managing phytopathogenic fungi (Basit *et al.*, 2021), little exploration has been done regarding their use in managing viruses and viroids. This exploration aims to reduce the impact of agrochemicals on vector control since, by nature, the best management involves implementing resistant varieties and preventive measures. As a result, the objective of this review is to explore the potential of PGPR for managing viruses and viroids causing diseases in tomatoes, along with examining the current research trends in this field within the country.

### **Viruses in tomato**

Viruses are infectious agents (obligate parasites) composed of nucleic acid (RNA and DNA) enveloped in proteins, capable of replicating within living cells (Hancinský *et al.*, 2020). They are exclusively dependent on the host cell's machinery for replication (Sastry, 2013b). Viruses can exhibit various shapes such as rigid and flexible rods, rigid rods, bacilli, polyhedra, geminate, among others (ICTV, 2021).

In 2011, the ten most significant viruses of international importance were identified by the global plant virology community. These included *Tobacco mosaic virus* (TMV), *Tomato spotted wilt virus* (TSWV), *Tomato yellow leaf curl virus* (TYLCV), *Cucumber mosaic virus* (CMV), *Potato virus Y* (PVY), *Cauliflower mosaic virus* (CaMV), *African cassava mosaic virus* (ACMV), *Plum pox virus* (PPV), *Brome mosaic virus* (BMV), and *Potato virus X* (PVX). Notably, six of these viruses infect tomatoes (Scholthof *et al.*, 2011). Worldwide, the key viruses causing losses in tomato cultivation comprise *Tomato leaf curl virus* (TYLCV), *Tobacco mosaic virus*, *Beet curly top virus* (BCTV), *Tomato bushy stunt virus* (TBSV), and *Tomato spotted wilt virus* (Sastry, 2013a). Since 2018, the emergence of *Tomato brown rugose fruit virus* (TBRFV) poses a latent threat due to the lack of tolerant varieties and its distribution in major production areas in Mexico (Cambrón-Crisantos *et al.*, 2019).

The International Committee on Taxonomy of Viruses (ICTV) has classified viruses into 189 families, 2,224 genera, and 9,110 species (<https://talk.ictvonline>).

org/taxonomy/) (ICTV, 2021). In tomatoes, approximately 312 viruses have been reported (Rivarez *et al.*, 2021), with more than 28 of them documented in Mexico (Table 1). Among the viruses reported in Mexico, ToBRFV, TYLCV, TSWV, *Pepino mosaic virus* (PepMV), and *Tomato wilt virus* (ToMarV) are of greater significance in tomatoes due to their economic impact and prevalence in both field and greenhouse conditions (García-Estrada *et al.*, 2022).

Among the primary viruses affecting tomatoes, over 28 viruses belong to the Begomovirus genus (ssDNA) (Hogenhout *et al.*, 2008), while the remaining viruses fall into 18 genera. Notably, *Orthotospovirus*, *Potyvirus*, *Tobamovirus*, and

**Table 1.** Main viruses reported in tomato (*Solanum lycopersicum*) in Mexico and the world.

Family / Genus / Species	Type genome	Reference
<b>Geminiviridae: Begomovirus</b> *Pepper golden mosaic virus (PepGMV), *Tomato chino La Paz virus (ToChLPV), *Tomato golden mottle virus (ToGMoV), Tomato leaf curl Bangladesh virus (ToLCBV), *Tomato mottle virus (ToMoV), *Tomato severe leaf curl virus (ToSLCV), *Tomato yellow leaf curl virus (TYLCV), Tobacco leaf curl virus (TLCV), Tomato bright yellow mottle virus (TBYMV), Tomato enation leaf curl virus (ToELCV), Tomato common mosaic virus (ToCmMV), Tomato curly stunt virus (ToCSV), Tomato chlorotic leaf distortion virus (ToCILDV), Tomato chlorotic mottle Guyane virus (ToCMoGV), Tomato dwarf leaf virus (ToDLV), Tomato golden mosaic virus (TGMV), Tomato golden vein virus (TGVV), Tomato interveinal chlorosis virus (ToICV), Tomato leaf curl New Delhi virus (ToLCNDV), Tomato leaf curl Palampur virus (ToLCPaV), Tomato leaf curl purple vein virus (ToLCPVV), Tomato mosaic Havana virus (ToMHaV), Tomato mottle leaf curl virus (TOMLCV), Tomato rugose yellow leaf curl virus (TRYLCV), Tomato yellow leaf curl Sardinia virus (TYLCSV), Tomato yellow mosaic virus (ToYMV), Tomato leaf curl virus Arusha virus (ToLCArV), Tomato leaf curl Ghana virus (ToLCGV), *Sinaloa tomato leaf curl virus (SToLCV), *Tomato leaf curl Sinaloa virus (ToLCSinV), <i>Lisianthus enation leaf curl virus</i> (LELCV), *Chino del tomate virus (CdTV), *Pepper huasteco yellow vein virus (PHYVV), *Squash leaf curl virus (SLCV), *Tomato apex necrosis virus (ToANV)	Bipartita, monopartita circular / ssDNA	Green <i>et al.</i> , 2007; Holguin-Peña <i>et al.</i> , 2004; Honguin-Peña <i>et al.</i> , 2007; Mauricio-Castillo <i>et al.</i> , 2007; CABI, 2020; Mauricio-Castillo <i>et al.</i> , 2007; Cardenas-Conejo <i>et al.</i> , 2010; Avedi <i>et al.</i> , 2021; Idris y Brown, 2007; Rojas <i>et al.</i> , 2005; Taniguchi <i>et al.</i> , 2023; Idris <i>et al.</i> , 1999; Lugo <i>et al.</i> , 2011; Barajas-Ortiz <i>et al.</i> , 2013; Zuñiga-Romano <i>et al.</i> , 2019
<b>Potyviridae: Potyvirus</b> *Tobacco etch virus (TEV), Potato virus Y (PVY), Pepper vein mottle virus (PVMV), Peru tomato mosaic virus (PTMV); *Tomato necrotic stunt virus (ToNSV)	Monopartita lineal / ssRNA +	CABI, 2020; Fernandez-Northcote y Fulton, 1980; Li <i>et al.</i> , 2012
<b>Amalgaviridae: Amalgavirus</b> *Southern tomato virus (SToV)	Lineal / dsRNA	Sabanadzovic <i>et al.</i> , 2009
<b>Virgoviridae: Tobamovirus</b> *Tobacco mosaic virus (ToMV), *Tomato brown rugose fruit virus (ToBRFV), *Tomato mottle mosaic virus (ToMMV), *Tomato mosaic virus (ToMV)	Monopartita lineal ssRNA +	CABI, 2020; Cambrón-Crisantos <i>et al.</i> , 2019; Zuñiga-Romano <i>et al.</i> , 2019


Table 1. Continue...

Family / Genus / Species	Type genome	Reference
<b>Secoviridae: <i>Nepovirus</i></b> <i>*Tobacco ringspot virus</i> (TRSV), <i>Tomato black ring virus</i> (TBRV)	Bipartita lineal / ssRNA +	CABI, 2020; Perez-Moreno <i>et al.</i> , 2004
<b>Secoviridae: <i>Torradovirus</i></b> <i>*Tomato torrado virus</i> (ToTV); <i>Tomato marchitez virus</i> (ToMarV), <i>*Tomato chocolate spot virus</i> (ToChSV), <i>Tomato chocolate virus</i> (ToChV)	Bipartita lineal / ssRNA +	CABI, 2020; Verbeek <i>et al.</i> , 2008
<b>Tombusviridae: <i>Tombusvirus</i></b> <i>*Tomato bushy stunt virus</i> (TBSV)	Monopartita lineal / ssRNA +	CABI, 2020; De la Torre-Almaráz <i>et al.</i> , 2004
<b>Closteroviridae: <i>Crinivirus</i></b> <i>*Tomato chlorosis virus</i> (ToCV), <i>*Tomato infectious chlorosis virus</i> (TICV)	Bipartita lineal / ssRNA +	CABI, 2020; Méndez-Lozano <i>et al.</i> , 2012
<b>Tospoviridae: <i>Orthospovirus</i></b> <i>*Tomato spotted wilt virus</i> (TSWV), <i>Chrysanthemum stem necrosis virus</i> (CSNV), <i>Pepper necrotic spot virus</i> (PNSV), <i>Tomato chlorotic spot virus</i> (TCSV), <i>Tomato necrotic ring virus</i> (TNRV) <i>Groundnut bud necrosis virus</i> (GBNV), <i>*Impatiens necrotic spot virus</i> (INSV)	ssRNA +	CABI, 2020; Nagata y de Ávila, 2000; Suganyadevi <i>et al.</i> , 2018; Zuñiga-Romano <i>et al.</i> , 2019; Honguin-Peña <i>et al.</i> , 2007
<b>Betaflexiviridae: <i>Carlavirus</i></b> <i>Cowpea mild mottle virus</i> (CPMMV)	Lineal / ssRNA +	EPPO, 2023
<b>Alphaflexiviridae: <i>Potexvirus</i></b> <i>*Pepino mosaic virus</i> (PepMV), <i>Potato virus X</i> (PVX)	Lineal / ssRNA +	EPPO, 2023; Zuñiga-Romano <i>et al.</i> , 2019
<b>Tymoviridae: <i>Tymovirus</i></b> <i>Eggplant mosaic virus</i> (EMV)	Lineal / ssRNA +	EPPO, 2023
<b>Secoviridae: <i>Fabavirus</i></b> <i>Broad bean wilt virus</i> (BBWV)	Bipartita lineal / ssRNA +	CABI, 2020
<b>Bromoviridae: <i>Alfamovirus</i></b> <i>*Alfalfa mosaic virus</i> (AMV)	Tripartita lineal / ssRNA	De la Torre-Almaráz <i>et al.</i> , 2003
<b>Bromoviridae: <i>Cucumovirus</i></b> <i>*Cucumber mosaic virus</i> (CMV), <i>Peanut stunt virus</i> (PSV), <i>Tomato aspermy virus</i> (TAV)	Tripartita lineal / ssRNA +	EPPO, 2023; Lecoq y Desbiez, 2012; Zuñiga-Romano <i>et al.</i> , 2019
<b>Bromoviridae: <i>Iilarvirus</i></b> <i>Tobacco streak virus</i> (TSV), <i>Parietaria mottle virus</i> (PMoV)	Tripartita lineal / ssRNA +	CABI, 2020
<b>Geminiviridae: <i>Curtovirus</i></b> <i>Beet curly top virus</i> (BCTV)	Monopartita circular / ssDNA	Chen y Gilbertson, 2016

\*Viruses reported in Mexico.

*Cucumovirus* exhibit a higher number of species following the *Begomovirus* genus. In some instances, viruses are transmitted by insect vectors (Figure 2); for instance, Potyviruses are transmitted by aphids, Orthotospoviruses by thrips (Aphidinae) (Gibbs *et al.*, 2008; Sastry, 2013b), and Begomoviruses by whiteflies (*Bemisia tabaci*, *Trialeurodes vaporariorum*) (Hogenhout *et al.*, 2008). This vector-mediated transmission adds complexity to the management of these phytopathogens.

Typically, symptoms induced by viruses are challenging to identify due to their occurrence in mixed viral infections (Sastry, 2013a). It is evident that these symptoms diminish the vigor, quality, and yield of crops (Figure 3) (Sastry, 2013b).



**Figure 2.** Main forms of transmission of viruses and viroids in tomato. A) Transmission by seed; B) Mechanical transmission, due to the use of work tools, manipulation of plants; C) Transmission by insect vectors.





**Figure 3.** Symptoms associated with viruses and viroids in tomatoes. A and B) Mosaic, leaf reduction, and mild to severe leaf distortion associated with *Tomato brown rugose fruit virus*; C and D) Stunting, fruit deformation, and purple discoloration in leaves caused by *Mexican papita viroid*; E) Yellow mosaic symptom associated with *Pepino mosaic virus*; F) Symptoms of stunting, deformation, and severe mosaic associated with *Begomovirus*; G and H) Symptoms of concentric rings and slight fruit deformation associated with *Tomato spotted wilt virus*; I) Mosaic in leaves caused by *Tobacco mosaic virus*.

Diseases caused by these phytopathogens pose a significant challenge to agriculture. Moreover, imminent climate change and agricultural practices have favored outbreaks of diseases, leading to their spread to unreported or previously unaffected areas (Jones and Naidu, 2019).

### **Viroids in tomato**

Viroids are infectious agents with lower structural and genetic complexity, consisting of a covalently closed circular RNA strand of low molecular weight (246 to 401 nucleotides in length) (Flores *et al.*, 1998). Some are pathogenic, while others replicate without inducing symptoms in the host (Flores *et al.*, 1998). Most viroids were initially discovered due to the damage they caused in various economically important crops (Flores and Randles, 2003).

The International Committee on Taxonomy of Viruses (ICTV) has classified 33 viroids into two families based on the cellular location of replication: Pospiviroidae (28 species) and Avsunviroidae (five species) (Di Serio and Flores, 2008). Each family consists of genera, and these genera comprise species with numerous sequence variants known as strains (Flores *et al.*, 1998; 2000). Upon entering the host cell, Pospiviroidae move to the nucleus before initiating replication, whereas Avsunsoviroidae replicate in the chloroplasts (Sastry, 2013b).

Since 1988, diseases naturally infected by viroids have been detected in several countries; all of the family Pospiviroidae, such as *Tomato planta macho viroid* (TPMVd) (Galindo *et al.*, 1982), *Mexican papita viroid* (MPVd) which could be considered as a variant of the TPMVd (Verhoeven *et al.*, 2011), *Potato spindle tuber viroid* (PSTVd) (Puchta *et al.*, 1990), *Tomato apical stunt viroid* (TASVd) (Walter *et al.*, 1980; Walter, 1987), *Citrus exocortis viroid* (CEVd) (Mishra *et al.*, 1991; Fagoaga y Duran-Vila, 1996; Verhoeven *et al.*, 2004), *Indian tomato bunchy top viroid* (Mishra *et al.*, 1991) which is considered a CEVd strain (Singh *et al.*, 2003), *Tomato chlorotic dwarf viroid* (TCDVd) (Singh *et al.*, 1999), and *Columnnea latent viroid* (CLVd) (Verhoeven *et al.*, 2004). In artificial infections, the tomato was successfully infected with *Chrysanthemum stunt viroid* (CSVd) (Matsushita y Kumar, 2009) and *Pepper chat fruit viroid* (PCFVd) (Table 2) (Verhoeven *et al.*, 2009).

All viroids are transmitted through mechanical inoculation (Sastry, 2013b) and by contact (Figure 2); for instance, through tools used in pruning, via clothing, through crop manipulation, and direct contact between nearby plants (Hammond, 2017). In other cases, transmission occurs through seeds, vegetative propagation, grafting, pollen, and insects (Figure 2) (Verhoeven *et al.*, 2004). The efficiency of viroids in being transmitted through seeds varies among plant and viroid species (Flores *et al.*, 2011; Singh *et al.*, 2003; Chung and Choi, 2008). Viroid infections



**Table 2.** Viroids that affect tomato (*Solanum lycopersicum*) in the world.

Species	Found	Symptoms in tomato*	Type of transmission	Losses and incidence	Reference
<i>Tomato planta macho viroid</i> = <i>Mexican papita viroid</i> (TPMVd=MPVd)	México (1982, 2008 and 2020) Canada (2008)	Stunting; chlorosis in the apex, tan or purple leaves, epinasty and severe deformation, reduction in the size and number of fruits and abortion of flowers.	Probable vector <i>Myzus persicae</i>  Seed by contaminated pollen	Causes significant losses in yield. Incidences of 5% on the crop.	Galindo <i>et al.</i> , 1982, 1986; Ling y Zhang, 2009; Mejorada-Cuellar, 2020; Ling y Bledsoe, 2009; Yanagisawa y Matsushita, 2017; Yanagisawa and Matsushita, 2018; Aviña-Padilla <i>et al.</i> , 2018; Matthews-Berry, 2010; Li y Ling, 2012
<i>Potato spindle tuber viroid</i> (PSTVd)	Discovered in 1922 distributed worldwide but eradicated in most countries. Australia (2011)	Epinasty, chlorosis, violet color with bright yellow central nervation, deformation and reduction in size, brittle, and necrotic central nervation, short internodes, with stunted shoots and necrosis, abortion of flowers, small and hard fruits with dark green spots and irregular maturing	Seed  Vector <i>M. persicae</i> (Transcap-sidación with <i>Potato leaf roll virus</i> )	Reduction of crop biomass and yield by up to 89%.  Incidence of 3% in cherry tomato ( <i>S. lycopersicum</i> cv. Perino).	Owens, 2007; Matthews-Berry, 2010; van Brunshot <i>et al.</i> , 2014; Yanagisawa y Matsushita, 2017; Matsushita <i>et al.</i> , 2011; Diener, 1987; Singh <i>et al.</i> , 2003 NSW Government, 2012; Mackie <i>et al.</i> , 2019
<i>Tomato apical stunt viroid</i> (TASVd)	Israel (1999 y 2000), Some countries in Asia and Africa, several European countries.	Short internodes; deformed, brittle leaves with yellowing; reduction and colorlessness in fruits (pale reddish)		Incidence of almost 100%	Antignus <i>et al.</i> , 2002, 2007; CABI, 2018; Matsuura <i>et al.</i> , 2010; Nielsen <i>et al.</i> , 2012
<i>Citrus exocortis viroid</i> (CEVd)	Distributed worldwide, mainly in citrus fruits	Chlorosis, tanning and distortion, reduction in growth.	Found in commercial seed lots	No reports	Mishra <i>et al.</i> 1991; Matthews-Berry, 2010; Verhoeven <i>et al.</i> , 2004 Constable <i>et al.</i> , 2019
<i>Tomato chlorotic dwarf viroid</i> (TCDVd)	Canada, U.S.A., Mexico, Japan, Hawaii (2017)	Reduction in growth and yellowing (chlorosis) of shoots, deformation of fruits, chlorosis turns bronze-colored, reddish and /or purple, epinasty Death of plants.	By active pollination by bumblebees ( <i>Bombus terrestris</i> and <i>B. ignites</i> )	100% incidence in greenhouses	Matsushita <i>et al.</i> , 2008; Singh <i>et al.</i> , 1999; Ling y Zhang, 2009; Verhoeven <i>et al.</i> , 2004 y 2007; Singh y Teixeira, 2006; Olmedo-Velarde <i>et al.</i> , 2018; Nie, 2012; Matthews-Berry, 2010; Antignus <i>et al.</i> , 2007; Matsuura <i>et al.</i> , 2010; Singh y Dilworth, 2009

Table 2. Continue...

Especie	Found	Symptoms in tomato*	Type of transmission	Losses and incidence	Reference
<i>Columnea latent viroid</i> (CLVd)	Netherlands and Belgium  In 2007 in Great Britain and France and in 2011 in Mali.	Severe deformation, tanning and burning of leaves, epinasty, chlorosis and necrosis in nervation, reduction in size or delay in development.	Found in commercial seed lots	Decline in production in Great Britain. Incidence on field of approximately 1.5%.	Nixon et al., 2010; Batuman y Gilbertson, 2013; Matthews-Berry, 2010; Steyer et al., 2010; Constable et al., 2019.
<i>Pepper chat fruit viroid</i> (PCFVd)	In 2009 in bell pepper plants and in 2013 in tomato seed shipments	Necrosis in young leaves and necrosis in nervation and petioles, stunting, reduction in fruit size.	Found in commercial seed lots	No reports	Verhoeven et al., 2009; Chambers et al., 2013; Constable et al., 2019

\*The described symptoms that each viroid can cause can be expressed together or separately in an infected plant.

in commercial tomato crops have been linked to the importation of seeds and ornamental plants (Batuman and Gilbertson, 2013; Van Brunschot *et al.*, 2014; Verhoeven *et al.*, 2012). Due to the diversity of viroid species, including some endemic ones, and the presence of mixed infections in tomatoes, Mexico represents a center of origin for viroids (Aviña-Padilla *et al.*, 2022).

Symptoms induced by viroids in tomatoes depend on the viroid species, cultivar, temperature, and light conditions. Plant responses are additionally influenced by RNA silencing, which plays a significant role in symptom development, combined with structural elements of viroids (Di Serio *et al.*, 2013; Flores *et al.*, 2015). The variability in symptoms is caused by different forms of gene expression. Moreover, viroid infections may or may not induce symptoms, but they generally lead to chlorosis, bronzing, leaf distortion, stunting, vein clearing, and discoloration, as well as malformation of flowers and fruits, reduced yield, and non-commercial fruit (Figure 1 C and D) (Singh *et al.*, 2003; Kovalskaya and Hammond, 2014).

### Management of diseases caused by viruses and viroids

There is a global trend towards the consumption of pesticide-free products. Regardless of the agricultural system and phytopathogen, it is ideal to adopt an Integrated Pest Management (IPM) approach that includes measures before, during, and after crop growth. As a first step, the use of virus-resistant cultivars is recommended (Rojas *et al.*, 2018), along with cultural sanitation practices as disease management measures (Karthika *et al.*, 2020).

The effective control of viruses and viroids involves early detection (diagnosis), eradication, and cultural control methods (Kovalskaya and Hammond, 2014).

Implementing appropriate control measures will help prevent or reduce the severity of viruses and/or viroids in plants. Therefore, diagnosis for virus identification is crucial for implementing management strategies. Diagnostic techniques may include ELISA, immunochromatography or strip tests, nucleic acid hybridization, polymerase chain reaction (PCR), and next-generation sequencers (González-Garza, 2017).

Given the significance of viral and viroidal diseases, alternative approaches have been sought, such as the use of beneficial microorganisms to protect the crop and enhance plant growth and productivity (Sofy *et al.*, 2019).

As mentioned earlier, the use of tolerant varieties has been implemented as part of the management virus and/or viroidal variety; however, in recent years, there has been a growing emphasis on issues related to the induction of natural defenses in plants (Ryals *et al.*, 1994; Kloepper *et al.*, 2004). The demand for biological products for pest and disease control, along with the use of inducing molecules (elicitors) capable of triggering defense responses in plants, is on the rise (Nasir *et al.*, 2014). It should be noted that the magnitude and effectiveness of the induced response will depend on the type of molecule, the signal, or its ability to induce secondary signaling within the tissue (Eder and Cosio, 1994).

By nature, plants possess the ability to defend themselves against phytopathogens through the production of substances that prevent or reduce damage caused by microorganisms. This response is triggered by plant-pathogen recognition after a local infection, where plants activate defense responses systemically to increase the magnitude and speed of the response against the pathogen (Delgado-Oramas, 2020). Induced resistance is a state where plants enhance their defenses against phytopathogen attacks and are triggered by the stimulation of chemical and/or biological inducers (Choudhary *et al.*, 2007). In general, induced resistance can function for the control of a broad spectrum of phytopathogens (Kloepper, 1993). Two types of pathogen-induced resistance are known in plants: Induced Systemic Resistance (ISR) and Systemic Acquired Resistance (SAR) (Camarena-Gutiérrez and Torre-Almaráz, 2007).

**Induced Systemic Resistance (ISR).** The activation of ISR requires the signaling of salicylic acid (SA), jasmonic acid (JA), and ethylene (ET) (Beneduzi *et al.*, 2012; Pieterse *et al.*, 2014). These products coordinately transduce extracellular stimuli recognized by host cell receptors into a large number of target molecules that integrate specific intracellular responses to external stimuli (Walters, 2009; Delgado-Oramas, 2020; Jankiewicz and Koltonowicz, 2012; Choi *et al.*, 2014). ISR has been observed in various plant species as part of defense reactions against fungi, bacteria, and viruses (Camarena-Gutiérrez and de la Torre-Almaráz, 2007).

**Signaling pathways mediated by ET, AJ, AS and their interaction.** The signaling pathways mediated by SA, JA, and ET do not operate independently (Maldonado-Cruz *et al.*, 2008; Derksen *et al.*, 2013). Through a cascade of signals from mitogen-activated protein kinases (MAPKs), the activation of ethylene-insensitive genes 1, 2, and 3 (EIN1, EIN2, and EIN3) occurs, regulating gene expression in response to pathogens and injuries (Zhu *et al.*, 2011). In a study on mustard (*Sinapis alba*), it was demonstrated that ethylene, in interaction with the mitochondrial alternative oxidase pathway (AOX), induces systemic resistance to Turnip mosaic virus (TuMV). This is attributed to the potential limitation of the virus's systemic infection and its accumulation in the plant (Zhu *et al.*, 2011).

Jasmonate signaling plays a crucial role in the induction of defense genes in plants against pests (Garnica *et al.*, 2012). The gene for coronatine-insensitive protein (COI1) serves as a central regulator in the JA signaling pathway. Upon binding with Jasmonoyl-Isoleucine (JA-Ile), it facilitates the release of the transcription factor MYC2, leading to the transcription of genes responsive to jasmonates (Garnica *et al.*, 2012). Evidence suggests that the expression of the COI1 gene and the N gene in the tobacco-TMV interaction imparts resistance to TMV (Liu *et al.*, 2004).

Salicylic acid (SA) is a pivotal signaling molecule in intracellular signal transduction. In addition to contributing to the release of H<sub>2</sub>O<sub>2</sub> and its active oxygen derivatives, it can induce the expression of genes related to defense (Shirasu *et al.*, 1997).

**Induced Systemic Response in tomato against phytopathogenic viruses mediated by PGPR.** The need to ensure crop quality and high yields with low environmental impact has led to the emergence of different inputs formulated with microorganisms, among which the use of Plant Growth-Promoting Rhizobacteria (PGPR) can be highlighted (González *et al.*, 2018; Canchignia *et al.*, 2015). These microorganisms enhance Systemic Resistance Induction (SRI) and are saprophytic bacteria that live freely in the rhizosphere. By colonizing the plant root system, they contribute through secretions, vitamins, hormones, and other growth factors that help improve plant growth and productivity (Walters, 2009). What is interesting is their use of the CVRs on foliage and seeds, observing different benefits (Table 3, Figure 4).

Most bacteria reported as Plant Growth-Promoting Rhizobacteria (PGPR) belong to the genera *Pseudomonas* and *Bacillus*. Additionally, bacteria from the following genera have been reported: *Aeromonas*, *Agrobacterium*, *Arthrobacter*, *Alcaligenes*, *Azospirillum*, *Azoarcus*, *Azotobacter*, *Burkholderia*, *Bradyrhizobium*, *Comamonas*, *Cyanobacteria*, *Enterobacter*, *Gluconacetobacter*, *Pizobielonas*, *Serratia*, *Variovorax*, *Streptomyces*, and *Xanthomonas* (Vessey, 2003).



**Table 3.** Plant growth-promoting rhizobacteria species used for virus management in tomato.

Viruses	PGPR	Result obtained	Reference
Cucumber mosaic virus (CMV)	<i>Pseudomonas fluorescens</i> , <i>Serratia marcescens</i>	Reduction in the area under the disease progress curve in diseased plants treated.	Raupach <i>et al.</i> , 1996
	<i>Bacillus amyloliquefaciens</i> (IN937a), <i>B. subtilis</i> (IN937b), <i>B. pumilus</i> (SE34)	Symptom reduction from 38% to 58% with PGPR, compared with 88% to 98% without PGPR.	Zehnder <i>et al.</i> , 2000, 2001
	<i>B. pumilus</i> (SE34), <i>B. amyloliquefaciens</i> (IN937a), <i>B. subtilis</i> (IN937b), <i>B. pumilus</i> (INR7, T4)	Infection rate from 27.5 (SE34) to 85% (T4), compared with control (87.5%). It favored the growth of plants and protected them against the virus.	Murphy <i>et al.</i> , 2003
	<i>Azospirillum lipoferum</i> (MRB16), <i>A. brasiliense</i> (SP7), <i>A. brasiliense</i> (N040), <i>Anabena oryzae</i>	Infected and treated plants increased their greenhouse yield to 48% and field yield to 40%.	Dashti <i>et al.</i> , 2007
	<i>P. aeruginosa</i> , <i>Stenotrophomonas rhizophilia</i>	Promoted vegetative growth and yield. Prevented infection in 91% of plants.	Dashti <i>et al.</i> , 2012
Tobacco mosaic virus (TMV)	<i>Pseudomonas</i> spp. (B-25)	Promoted plant growth and yield. Reduced the incidence of the virus in plants.	Kirankumar <i>et al.</i> , 2008
	<i>Bacillus</i> spp. <i>Bacillus amyloliquefaciens</i> (TBorg1)	Favored the development of roots in infected tobacco plants. 90 % reduction in virus buildup. Increased total soluble carbohydrates, proteins and ascorbic acid.	Wang <i>et al.</i> , 2009; Choi <i>et al.</i> , 2014 Abdelkhalek <i>et al.</i> , 2022
	<i>P. fluorescens</i> SM90 y <i>B. subtilis</i> DR06	Increased defense-related genes NPR1, COI1 AND PR1-a, promotes growth in plants infected with the virus.	Sharaf <i>et al.</i> , 2023
	<i>B. amyloliquefaciens</i> (IN937a) <i>B. pumilus</i> (SE34)+ <i>B. amyloliquefaciens</i> (IN937a) <i>B. pumilus</i> (SE34)+ <i>B. sphaericus</i> (SE56)+ <i>B. amyloliquefaciens</i> (IN937a)	It reduced severity in the field and was the treatment that best controlled the disease.	Abdalla <i>et al.</i> , 2017
Tomato mottle virus (ToMoV)	<i>B. amyloliquefaciens</i> (IN 937a) <i>B. subtilis</i> (IN 937b) <i>B. pumilus</i> (SE34)	Reduced the incidence of the virus in plants by 30%.	Murphy <i>et al.</i> , 2000; Zehnder <i>et al.</i> , 2001
		Reduced the development of symptoms and incidence.	

Table 3. Continue...

Viruses	PGPR	Result obtained	Reference
Tomato spotted wilt virus (TSWV)	<i>P. fluorescens</i> (CHA0)	Increased yield and growth of infected and treated plants. Reduced the viral load.	Kandan <i>et al.</i> 2005; 2002
	<i>P. fluorescens</i> (CHA0+CoT-1)		
	<i>P. fluorescens</i> (CHA0+CoT-1+CoP-1)		
Tomato yellow leaf curl virus (TYLCV)	<i>B. amyloliquefaciens</i> (MBI600)	Reduced the incidence of the virus by 80%.	Beris <i>et al.</i> , 2018
Potato virus Y	<i>Enterobacter asburiae</i> (BQ9)	Reduced the severity in treated and infected plants by 52%.	Li <i>et al.</i> , 2016
Groundnut bud necrosis virus (GBNV)	<i>Bacillus amyloliquefaciens</i> (strain MBI600)	Reduced the systemic accumulation of viruses.	Beris <i>et al.</i> , 2018
	<i>Bacillus amyloliquefaciens</i> (VB7)	Reduced symptoms by around 84%.	Vanthana <i>et al.</i> , 2019




Figure 4. Forms of application and mechanisms of action of Plant Growth-Promoting Rhizobacteria (PGPR) used to protect tomatoes from viral infections.

There are studies on the use of PGPR employed as microbial inoculants to protect tomato plants against viral infections, where, additionally, an increase in crop yield was recorded (Table 3). For instance, in Mexico, Samaniego *et al.* (2017) and Hernández-Santiago *et al.* (2020) reported that the application of *Bacillus* to infected tomato plants not only led to an increase in plant height and weight but also induced systemic resistance. In recent years, research on Systemic Resistance Induction (SRI) using PGPR against economically relevant viruses (20 viruses) in various crops has gained significance (Sofy *et al.*, 2019). However, despite efforts to counteract viruses, research remains scarce, particularly in the case of viroids. The work has mainly focused on crops such as *Musa* spp., *Vigna unguiculata*, *V. mungo*, *Vicia faba*, *Momordica charantia*, *Cucumis sativus*, *Citrullus lanatus*, *Solanum lycopersicum*, *S. tuberosum*, *Capsicum annuum*, *Nicotiana tabacum*, *N. tabacum* cv. *Xanthi-nc*, *Arabidopsis thaliana*, *Chenopodium quinoa*, *Helianthus annuus*, *Phaseolus vulgaris*, *Datura metel*, and *Cucurbita maxima* (Sofy *et al.*, 2019).

There are other microorganisms, such as endophytic fungi (*Trichoderma* spp.), which are also resistance inducers in plants, and they are even sold as biocontrollers of phytopathogenic fungi by companies in Mexico, to be applied to various crops. However, the use of these microorganisms and their effect on infected plants by viruses and viroids has been shallowly studied. Nevertheless, some species of rhizosphere and endophytic fungi seem to have a favorable effect as plant growth promoters in virus-infected tomato plants (Ramos-Villanueva *et al.*, 2023).

## CONCLUSIONS

Diseases caused by viruses and viroids impact the yield and quality of tomatoes, as well as other economically significant crops worldwide. In Mexico, there are records of over 28 viruses, mostly belonging to the Begomovirus genus. However, research on Plant Growth-Promoting Rhizobacteria (PGPR) in tomatoes as inducers of resistance against these pathogens or to enhance crop health is limited. The use of *B. subtilis* has been reported for controlling *Tobacco mosaic virus* in tomatoes (Samaniego *et al.*, 2017; Hernández-Santiago *et al.*, 2020). Globally, PGPR such as *Pseudomonas*, *Serratia*, *Bacillus*, *Azospirillum*, *Anabena*, *Enterobacter*, and *Stenotrophomona* have been evaluated against nine virus species in tomatoes, with a focus on *Cucumber mosaic virus*. Nevertheless, it is essential to strengthen such research due to the significance of tomato cultivation in Mexico. Promisingly, PGPR has demonstrated effects in reducing incidence and severity by up to 80%. Additionally, a yield increase of up to 48% has been recorded compared to plants without PGPR application. Therefore, the use of growth promoting microorganisms

are a promising alternative for integration into an integrated management of viral infections, given its compatibility with the environment and experimentally effective for the control of a variety of viruses.

## LITERATURE CITED

- Abdalla OA, Bibi S and Zhang S. 2017. Application of plant growth-promoting rhizobacteria to control *Papaya ringspot virus* and *Tomato chlorotic spot virus*. *Archives of Phytopathology and Plant Protection* 50: 584–597. <https://doi.org/10.1080/03235408.2017.1352248>
- Abdelkhalek A, Aseel DG, Király L, Künstler A, Moawad H and Al-Askar AA. 2022. Induction of systemic resistance to *Tobacco mosaic virus* in tomato through foliar application of *Bacillus amyloliquefaciens* Strain TBorg1 culture filtrate. *Viruses* 14: 1830. <https://doi.org/10.3390/v14081830>
- Antignus Y, Lachman O and Pearlsman M. 2007. The spread of Tomato apical stunt viroid (TASVd) in greenhouse tomato crops is associated with seed transmission and bumble bee activity. *Plant Disease* 91:47-50.
- Antignus Y, Lachman O, Pearlsman M, Gofman R and Bar-Joseph M. 2002. A new disease of greenhouse tomatoes in Israel caused by a distinct strain to *Tomato apical stunt viroid* (TASVd). *Phytoparasitica* 30: 502 - 510.
- Avedi EK, Adediji AO, Kilalo DC, Olubayo FM, Macharia I, Ateka EM, Machuka EM and Mutuku JM. 2021. Metagenomic analyses and genetic diversity of Tomato leaf curl Arusha virus affecting tomato plants in Kenya. *Virology journal* 18(1): 2. <https://doi.org/10.1186/s12985-020-01466-z>
- Aviña-Padilla K, Rivera-Bustamante R, Kovalskaya NY and Hammond WR. 2018. Pospiviroid Infection of Tomato Regulates the Expression of Genes Involved in Flower and Fruit Development. *Viruses* 10: 516. <https://doi.org/10.3390/v10100516>.
- Batuman O and Gilbertson RL. 2013. First Report of *Columnea latent viroid* (CLVd) in tomato in Mali. *Disease Note* 97 (5). <https://doi.org/10.1094/PDIS-10-12-0920-PDN>
- Barajas-Ortiz M, León-Sicairo CR, López-Valenzuela JA, Reyes-Moreno C, Valdez-Ortiz A, Velarde-Félix S, Peraza\_Garay F and Garzón-Tiznado JA. 2013. Transmission Efficiency of *Tomato Apex Necrosis Virus* by *Bemisia tabaci* (Hemiptera: Aleyrodidae) Biotype B in Tomato. 106(4): 1559-1565. <https://doi.org/10.1603/EC12223>
- Basit A, Shah ST, Muntha ST and Mohamed HI. 2021. Plant Growth-Promoting Rhizobacteria (PGPR) as Biocontrol Agents for Viral Protection. In: Mohamed, H.I., El-Beltagi, H.ED.S., Abd-Elsalam, K.A. (eds) *Plant Growth-Promoting Microbes for Sustainable Biotic and Abiotic Stress Management*. Springer, Cham. [https://doi.org/10.1007/978-3-030-66587-6\\_8](https://doi.org/10.1007/978-3-030-66587-6_8)
- Beneduzi A, Ambrosini A and Passaglia LMP. 2012. Plant growth-promoting rhizobacteria (PGPR): Their potential as antagonists and biocontrol agents. *Genetics and Molecular Biology* 35(4 Suppl):1044-1051.
- Beris D, Theologidis I, Skandalis N and Vassilakos N. 2018. *Bacillus amyloliquefaciens* strain MBI600 induces salicylic acid dependent resistance in tomato plants against *Tomato spotted wilt virus* and *Potato virus Y*. *Scientific Reports* 8:10320. <https://doi.org/10.1038/s41598-018-28677-3>
- CABI. 2020. Datasheet. Centre for Agricultural Bioscience International. <https://www.cabi.org/>
- CABI. 2018. Tomato apical stunt viroid. Datasheet. 20/04/18 Original text by: Rosemarie W Hammond, Molecular Plant Pathology Laboratory, USDA-ARS, Beltsville, Maryland, USA. <https://www.cabi.org/isc/datasheet/52804#todistribution>
- Camarena-Gutiérrez G y de la Torre-Almaráz R. 2007. Resistencia sistémica adquirida en plantas: Estado actual. *Revista Chapingo Serie Ciencias Forestales y del Ambiente* 13(2): 157-162.
- Cambrón-Crisantos JM, Rodríguez-Mendoza J, Valencia-Luna JB, Alcasio-Rangel S, García-Ávila CJ, López-Buenfil JA and Ochoa-Martínez DL. 2019. First report of *Tomato brown rugose fruit virus* (ToBRFV) in Michoacan, Mexico. *Mexican Journal of Phytopathology* 37(1): 185-192. <https://doi.org/10.18781/R.MEX.FIT.1810-5>
- Canchignia H, Cruz N, Barrera A, Morante-Cariel J, Canchignia M and Peñafiel M. 2015. Applying Plant Growth Promoting Rhizobacteria (PGPR) of the Genus *Pseudomonas* spp as Biological Controllers of Insects and Plague Nematodes. *Revista Ciencia y Tecnología* 8(1): 25-30. <https://dialnet.unirioja.es/servlet/articulo?codigo=5319281>



- Cardenas-Conejo Y, Arguello-Astorga G, Poghsyan A, Hernández-González J, Lebsky V, Holguin-Peña J, Medina-Hernández D and Vega-Peña S. 2010. First Report of *Tomato yellow leaf curl virus* Co-infecting Pepper with tomato chino La Paz virus in Baja California Sur, Mexico. *Plant Disease* 94(10): 1266. [10.1094/PDIS-06-10-0444](https://doi.org/10.1094/PDIS-06-10-0444)
- Constable F, Chambers G, Penrose L, Daly A, Mackie J, Davis K, Rodoni B and Gibbs M. 2019. Viroid-infected Tomato and Capsicum Seed Shipments to Australia *Viruses* 11: 98. <https://doi.org/10.3390/v11020098>.
- Chambers GA, Seyb AM, Mackie J, Constable FE, Rodoni BC, Letham D, Davis K and Gibbs MG. 2013. First Report of Pepper chat fruit viroid in Traded Tomato Seed, an Interception by Australian Biosecurity. *Disease notes* 97 (10). <https://doi.org/10.1094/PDIS-03-13-0293-PDN>
- Choi HK, Song GC, Yi HS and Ryu CM. 2014. Field Evaluation of the Bacterial volatile derivative 3-Pentanol in priming for induced resistance in Pepper. *Journal of Chemical Ecology* 40(8):882-92.
- Chen LFR, Gilvertson. 2016. Transmission of Curtoviruses (*Beet curly top virus*) by the Beet Leafhopper (*Circulifer tenellus*). In: Brown JK. (Eds.) Vector-mediated transmission of plant pathogens. APS <https://doi.org/10.1094/9780890545355.017>.
- Choudhary DK, Prakash A and Johri BN. 2007. Induced systemic resistance (ISR) in plants: Mechanism of action. *Indian Journal of Microbiology* 47(4): 289–297. <https://doi.org/10.1007/s12088-007-0054-2>
- Chung BN and Choi GS. 2008. Incidence of *Coleus blumei* viroid 1 in Seeds of Commercial Coleus in Korea. *The Plant Pathology Journal* 24(3): 305–308. <https://doi.org/10.5423/PPJ.2008.24.3.305>
- Dashti NH, Ali NY, Cherian VM and Montasser MS. 2012. Application of plant growth-promoting rhizobacteria (PGPR) in combination with a mild strain of *Cucumber mosaic virus* (CMV) associated with viral satellite RNAs to enhance growth and protection against a virulent strain of CMV in tomato. *Canadian Journal of Plant Pathology* 34:177–186. <https://doi.org/10.1080/07060661.2012.685495>
- Dashti NH, Montasser MS, Ali NY, Bhardwaj RG and Smith DL. 2007. Nitrogen biofixing bacteria compensate for the yield loss caused by viral satellite RNA associated with *Cucumber mosaic virus* in tomato. *The Plant Pathology Journal* 23:90–96. <https://doi.org/10.5423/ppj.2007.23.2.090>
- De la Torre-Almaráz R, Salazar-Segura M y Valverde RA. 2003. Etiología del moteado amarillo del tomar de cáscara (*Physalis ixocarpa* B.) en México. *Agrociencia* 37(3): 277-289. <https://www.redalyc.org/pdf/302/30237307.pdf>
- De la Torre Almaráz R, Monsalvo-Reyes AC, Méndez-Lozano J y Rivera-Bustamante RF. 2004. Caracterización de un nuevo geminivirus asociado con los síntomas de moteado amarillo de la okra (*Abelmoschus esculentus*) en México. *Agrociencia* 38(2): 227-238. <https://www.redalyc.org/pdf/302/30238210.pdf>
- Delgado-Oramas PB. 2020. La resistencia inducida como alternativa para el manejo de plagas en las plantas de cultivo. *Revista de Protección Vegetal*. 35(1). e07. Epub 01 de abril de 2020. [http://scielo.sld.cu/scielo.php?script=sci\\_arttext&pid=S1010-27522020000100001&lng=es&tlng=es](http://scielo.sld.cu/scielo.php?script=sci_arttext&pid=S1010-27522020000100001&lng=es&tlng=es).
- Derksen H, Rampitschb C and Daayf F. 2013. Signaling cross-talk in plant disease resistance. *Plant Science*, 207:79-87.
- Di Serio F and Flores F. 2008. Viroids: Molecular implements for dissecting RNA trafficking in plants. *RNA Biology* 5: 128 – 131. <https://doi.org/10.4161/rna.5.3.6638>
- Di Serio F, De Stradis A, Delgado S, Flores R and Navarro B. 2013. Cytopathic effects incited by viroid RNAs and putative underlying mechanisms. *Frontiers in Plant Science* 3:288. <https://doi.org/10.3389/fpls.2012.00288>
- Diener TO. 1987. Biological Properties. In: Diener T.O. (eds) *The Viroids. The Viruses*. Springer, Boston, MA. [https://doi.org/10.1007/978-1-4613-1855-2\\_2](https://doi.org/10.1007/978-1-4613-1855-2_2)
- Eder J and Cosio EG. 1994. Elicitors of plant defense responses. *International review of cytology* 148:1-36.
- Fagoaga C and Duran-Vila N. 1996. Naturally occurring variants of *citrus exocortis viroid* in vegetable crops. *Plant Pathology* 45: 45–53.
- FAO (Food and agriculture organization statistics). 2020. Consulted 16-December-2020. Disponible en: <http://faostat.fao.org/site/567/DesktopDefault.aspx?PageID=567#ancor>
- Fernandez-Northcote EN and Fulton RW. 1980. Detection and characterization of Peru tomato virus strains infecting pepper and tomato in Peru. *Phytopathology* 70:315-320. [https://www.apsnet.org/publications/phytopathology/backissues/Documents/1980Articles/Phyto70n04\\_315.PDF](https://www.apsnet.org/publications/phytopathology/backissues/Documents/1980Articles/Phyto70n04_315.PDF)

- Flores R and Randles J. 2003. Viroids: Properties, Detection, Diseases and Their Control Ahmed Hadidi, Science publishers, Inc. United States of America. 392 p.
- Flores R, Minoia S, Carbonell A, Gisel A, Delgado S, López-Carrasco A, Navarro B and Di Serio F. 2015. Viroids, the simplest RNA replicons: How they manipulate their hosts for being propagated and how their hosts react for containing the infection. *Virus Research* 209:136-145.
- Flores R, Randles JW, Bar-Joseph M and Diener TO. 1998. A proposed scheme for viroid classification and nomenclature. *Archives of Virology* 143: 623-629.
- Flores R, Randles JW, Bar-Joseph M and Diener T.O. 2000 Viroids. *In: Virus Taxonomy, 7<sup>th</sup> Report International of the Committee Taxonomy Viruses*. Regenmortel MHV, Fauquet CM, Bishop DHL, Carstens EB, Estes MK, Lemon SM, Maniloff J, Mayo MA, McGeoch DJ, Pringle CR and Wickner RB. (eds.), San Diego, Academic Press, 1009-1024 pp.
- Flores R, Di Serio F, Navarro B, Duran-Vila NURIA and Owens RA. 2011. Viroids and viroid diseases of plants. *Studies in Viral Ecology: Microbial and Botanical Host Systems* 1: 311-346.
- Galindo J, López M and Aguilar T. 1986. Significance of *Myzus persicae* in the spread of *tomato planta macho viroid*. *Fitopatología Brasil* 2: 400-410.
- Galindo J, Smith DR and Diener TO. 1982. Etiology of planta macho, a viroid disease of tomato. *Phytopatology* 72: 49–54.
- García-Estrada RS, Díaz-Lara A, Aguilar-Molina VH and Tovar-Pedraza JM. 2022. Viruses of economic impact on tomato crops in Mexico: From diagnosis to management-A review. *Viruses* 14:1251. <https://doi.org/10.3390/v14061251>
- Garnica VA, Raya GJ, López BJ y Beltrán PE. 2012. Señalización del ácido jasmónico y el óxido nítrico durante el desarrollo de *Arabidopsis thaliana*. *Ciencia Nicolaita* 55:28-41.
- Gibbs AJ, Ohshima K, Phillips MJ and Gibbs MJ. 2008. The Prehistory of Potyviruses: Their Initial Radiation Was during the Dawn of Agriculture. *PLOS ONE* 3(6): e2523. <https://doi.org/10.1371/journal.pone.0002523>
- González-Garza R. 2017. Evolución de técnicas de diagnóstico de virus fitopatógenos. *Revista Mexicana de Fitopatología* 35(3):591-610. <https://doi.org/10.18781/r.mex.fit.1706-1>.
- Green SK, Tsai WS, Shih SL, Black LL, Rezaian A, Rashid MH, Roff MMN, Myint YY and Hong LTA. 2007. Molecular Characterization of Begomoviruses Associated with Leafcurl Diseases of Tomato in Bangladesh, Laos, Malaysia, Myanmar, and Vietnam. *Disease Notes* 85(12). <https://doi.org/10.1094/PDIS.2001.85.12.1286A>
- Hanssen IM and Lapidot M. 2012. Major tomato viruses in the Mediterranean basin. *In: Loebenstein G and Lecoq H (Ed). Advances in Virus Research: Viruses and virus diseases of vegetables in the Mediterranean Basin* 84:31-66. <https://doi.org/10.1016/B978-0-12-394314-9.00002-6>.
- Hancinský R, Mihálik D, Mrkvová M, Candresse T and Glasa M. 2020. *Plants* 9 (667): 17 p. <https://doi.org/10.3390/plants9050667>
- Hammond WR. 2017. Viroids and Satellites. *Economic Significance of Viroids in Vegetable and Field Crops*. Chapter1. 5-13 pp. <http://dx.doi.org/10.1016/B978-0-12-801498-1.00001-2>.
- Hernández-Martínez J, García-Mata R, Valdivia-Alcalá y Omaña-Silvestre JM. 2004. Evolución de la competitividad y rentabilidad del cultivo del tomate rojo (*Lycopersicon esculentum* L.) en Sinaloa, México. *Agrociencia* 38: 431-436.
- Hernández-Santiago R, Vargas-Hernández M y Zamora-Macorra EJ. 2020. Evaluation of TMV resistance inducers in tomato. *Revista Mexicana Ciencias Agrícolas* 11(2): 377-390.
- Hogenhout SA, Ammar ED, Whitfield AE and Redinbaugh MG. 2008. Insect Vector Interactions with Persistently Transmitted Viruses. *Annual Review of Phytopathology* 46: 327–359. <https://doi.org/10.1146/annurev.phyto.022508.092135>
- Honguin-Peña RJ, Arguello-Astorga GR, Brown KJ and Rivera-Bustamante RF. 2007. A New Strain of Tomato chino La Paz virus associated with a leaf curl disease of tomato in Baja California Sur, Mexico. *Disease Notes* 90(7): 973. <https://doi.org/10.1094/PD-90-0973B>
- Holguín-Peña R. J., Vázquez-Juárez R. and Rivera-Bustamante R. F. 2004. *Pepper golden mosaic virus* affecting tomato crops in the Baja California Peninsula, Mexico. *Plant Disease* 88(2): 221. <https://doi.org/10.1094/PDIS.2004.88.2.221A>
- ICTV. 2021. International Committee on Taxonomy of Viruses. <https://talk.ictvonline.org>
- Idris AM and Brown JK. 2007. Sinaloa Tomato Leaf Curl Geminivirus: Biological and Molecular Evidence for a New Subgroup III Virus. *Virology*. <https://doi.org/10.1094/PHYTO.1998.88.7.648>

- Jankiewicz U and Koltonowicz M. 2012. The involvement of *Pseudomonas* bacteria in Induced Systemic Resistance in Plants (Review). *Applied Biochemistry and Microbiology* 48(39):244-249.
- Jones RAC and Naidu RA. 2019. Global Dimensions of Plant Virus Diseases: Current Status and Future Perspectives. *Annual Review of Virology* 6:387-409. <https://doi.org/10.1146/annurev-virology-092818-015606>
- Kandan A, Commare RR, Nandakumar R, Ramiah M, Raguchander T and Samiyappan R. 2002. Induction of phenylpropanoid metabolism by *Pseudomonas fluorescens* against tomato spotted wilt virus in tomato. *Folia Microbiologica* 47:121–129. <https://doi.org/10.1007/bf02817669>
- Kandan A, Ramiah M, Vasanthi VJ, Radjacommare R, Nandakumar R, Ramanathan A, Samiyappan R. 2005. Use of *Pseudomonas fluorescens*-based formulations for management of *Tomato spotted wilt virus* (TSWV) and enhanced yield in tomato. *Biocontrol Science and Technology* 15:553–569. <https://doi.org/10.1080/09583150500088546>
- Karthika S, Varghese S and Jisha MS. 2020. Exploring the efficacy of antagonistic rhizobacteria as native biocontrol agents against plant diseases. *3 Biotech* 10:320. <https://doi.org/10.1007/s13205-020-02306-1>
- Karthika S, Midhun SJ and Jisha MS. 2020. A potential antifungal and growth-promoting bacterium *Bacillus* sp. KTMA4 from tomato rhizosphere. *Microbial Pathogenesis* 8: 142: 104049. [10.1016/j.micpath.2020.104049](https://doi.org/10.1016/j.micpath.2020.104049)
- Kirankumar R, Jagadeesh KS, Krishnaraj PU and Patil MS. 2008. Enhanced growth promotion of tomato and nutrient uptake by plant growth-promoting rhizobacterial isolates in presence of *Tobacco mosaic virus* pathogen. *Karnataka Journal Agriculture Sciences* 21:309–311.
- Kloepper JW. 1993. Plant growth promoting rhizobacteria as *biological* control agents. In: Metting, F. B (ed) *Soil microbial ecology— applications in agricultural and environmental management*. Dekker, New York, pp 255–274.
- Kloepper JW, Ryu CM and Zhang S. 2004. Induced systemic resistance and promotion of plant growth by *Bacillus* spp. *Phytopathology* 94:1259–1266. <https://doi.org/10.1094/phyto.2004.94.11.1259>
- Kloepper JW, Tuzun S and Kuć JA. 1992. Proposed definitions related to induced disease resistance. *Biocontrol Science and Technology* 2:349-351. <https://doi.org/10.1080/09583159209355251>
- Lecoq H and Desbiez C. 2012. Viruses of Cucurbit Crops in the Mediterranean Region: An Ever-Changing Picture. In: Loebenstein G and Lecoq H (eds.) *Viruses and Virus Diseases of Vegetables in the Mediterranean Basin* 84: 67-126. <https://doi.org/10.1016/B978-0-12-394314-9.00003-8>
- Li H, Ding X, Wang Ch, Ke H, Wu Z, Wang Y, Liu and H Guo, J. 2016. Control of *Tomato yellow leaf curl virus* disease by *Enterobacter asburiae* BQ9 as a result of priming plant resistance in tomatoes. *Turkish Journal Biology*. 40:150–159. <https://doi.org/10.3906/biy-1502-12>
- Li, R. and Ling K. 2012. Molecular and biological characterization of *Mexican papita viroid*. *Tomato Disease Workshop*, 28 p.
- Li R, Gao S, Hernandez AG, Wechter WP, Fei Z and Ling KS. 2012. Deep Sequencing of Small RNAs in Tomato for Virus and Viroid Identification and Strain Differentiation. *Plos One*. <https://doi.org/10.1371/journal.pone.0037127>
- Ling, K. and M. Bledsoe. 2009. First Report of *Mexican Papita Viroid* Infecting Greenhouse Tomato in Canada. *Plant Disease*, 93: 839.
- Ling KS and Zhang W. 2009. First Report of a Natural Infection by *Mexican Papita Viroid* and *Tomato Chlorotic Dwarf Viroid* on Greenhouse Tomatoes in Mexico. *Plant Disease*, 93: 1126.
- Liu Y, Schiff M and Dinesh KSP. 2004. Involment of MEK1 MAPKK, NTF6 MAPK, WRKY/MYB transcription factors, COI1 and CTR1 in N-mediated resistance to tobacco mosaic virus. *The Plant Journal*.38(5):800-809.
- Lugo MOY, Guzmán UR, García ERS y León FJ. 2011. Geminivirus transmitidos por mosca blanca (*Bemisia tabaci*) en tomate del Valle Agrícola de Culiacán, Sinaloa. *Revista Mexicana de Fitopatología* 29:109-118. <https://www.scielo.org.mx/pdf/rmf/v29n2/v29n2a3.pdf>
- Mackie EA, Barbetti JM, Rodoni B, McKirdy SJ and Jones ACR. 2019. Effects of a *Potato Spindle Tuber Viroid* Tomato Strain on the Symptoms, Biomass, and Yields of Classical Indicator and Currently Grown Potato and Tomato Cultivars. *Plant disease*. <https://doi.org/10.1094/PDIS-02-19-0312-RE>
- Maldonado-Cruz E, Ochoa-Martínez DL y Tlapal-Bolaños B. 2008. Efecto del ácido acetil salicílico y *Bacillus subtilis* en la infección causada por *Cucumber mosaic virus* en calabacita. *Revista Chapingo Serie horticultura* 14(1):55-59.

- Matthews-Berry S. 2010. Emerging viroid threats to UK tomato production. The Food and Environment Research Agency (Fera). Plant disease Factsheet, 5 p.
- Matsura S, Matsushita Y, Kozuka Shimizu S and Tsuda S. 2010. Transmission of *Tomato chlorotic dwarf viroid* by bumblebees (*Bombus ignitus*) in tomato plants. *European Journal of Plant Pathology* 126: 111. <https://doi.org/10.1007/s10658-009-9515-2>
- Matsushita Y, Usugi T and Tsuda S. 2011. Distribution of *Tomato chlorotic dwarf viroid* in floral organs of tomato. *European Journal of Plant Pathology* 130(4): 441–447.
- Matsushita Y, Kanda A, Usugi T and Tsuda S. 2008. First report of a *Tomato chlorotic dwarf viroid* disease on tomato plants in Japan. *Journal of General Plant Pathology* 74: 182-184.
- Matsushita Y. and Kumar PKR. 2009. *In vitro* transcribed *Chrysanthemum stunt viroid* (CSVd) RNA is infectious to Chrysanthemum and other plants. *Phytopathology* 99: 58–66.
- Mauricio-Castillo JA, Argüello-Astorga GR, Ambriz-Granados S, Alpuche-Solís AG and Monreal-Vargas CT. 2007. First Report of *Tomato golden mottle virus* on *Lycopersicon esculentum* and *Solanum rostratum* in Mexico. *Plant Disease* 91(11): 1513. <https://doi.org/10.1094/PDIS-91-11-1513B>
- Mejorada-Cuellar BY, Zamora-Macorra EJ, Vilchis-Zimuta R and Zamora-Macorra M. 2020. Systemic acquired resistance to manage the *Mexican papita viroid* in tomato crop (*Solanum lycopersicum* L.). *Agroproductividad* 13(7): 75-80. <https://doi.org/10.32854/agrop.vi.1686>
- Méndez-Lozano J, Magallanes-Tapia MA, Romero-Romero JL, Camacho-Beltrán E, Orduño-Vega WL, Leyva-López NE, Santos-Cervantes ME and Félix-Gastélum R. 2012. Tomato infectious chlorosis virus Associated with Tomato Diseases in Baja California, Mexico. *Disease Notes* 96(8). <https://doi.org/10.1094/PDIS-02-12-0196-PDN>
- Mishra MD, Hammond RW, Owens RA, Smith DR and Diener TO. 1991. Indian bunchy top disease of tomato plants is caused by a distinct strain of *citrus exocortis viroid*. *Journal of General Virology* 72: 1781–1785.
- Murphy JF, Reddy MS, Ryu CM, Kloepper JW and Li R. 2003. Rhizobacteria-mediated growth promotion of tomato leads to protection against *Cucumber mosaic virus*. *Phytopathology* 93:1301–1307. <https://doi.org/10.1094/phyto.2003.93.10.1301>
- Murphy JF, Zehnder GW, Schuster DJ, Sikora EJ, Polston JE and Kloepper JW. 2000. Plant growth-promoting rhizobacterial mediated protection in tomato against *Tomato mottle virus*. *Plant Disease* 84:779–784. <https://doi.org/10.1094/pdis.2000.84.7.779>
- Nagata T and de Ávila AC. 2000. Transmission of Chrysanthemum Stem Necrosis Virus, a Recently Discovered Tospovirus, by Two Thrips Species. *Phytopathology* 148: 123-125. <https://doi.org/10.1046/j.1439-0434.2000.00475.x>
- Nasir MN, Polo LD, Luzuriaga LW, Deleu M, Lins L, Ongena M and Fauconnier ML. 2014. New alternatives to chemical pesticides: deciphering the action mechanisms of lipid-based plant elicitors via complementary biophysical and biological approaches. In: 19th National Symposium on Applied Biological Sciences. Gembloux, Bélgica. (<https://orbi.uliege.be/handle/2268/165417>).
- Nie X. 2012. Analysis of Sequence Polymorphism and Population Structure of Tomato chlorotic dwarf viroid and Potato spindle tuber viroid in Viroid-Infected Tomato Plants. *Viruses* 4(6):940-53. <https://doi.org/10.3390/v4060940>.
- Nielsen SL, Enkegaard A, Nicolaisen M, Kryger P, Viršček MM, Pleško IM, Kahrer A and Gottsberger RA. 2012. No transmission of *Potato spindle tuber viroid* shown in experiments with thrips (*Frankliniella occidentalis*, *Thrips tabaci*), honey bees (*Apis mellifera*) and bumblebees (*Bombus terrestris*). *European Journal Plant Pathology* 133: 505–509 <https://doi.org/10.1007/s10658-012-9937-0>
- Nixon T, Glover R, Matthews-Berry S, Daly ME, Hobden C, Lambourne V and Harju A. Skelton. 2010. *Columnnea latent viroid* (CLVd) in tomato: the first report in the United Kingdom. *Plant pathology* 59 (2): 392. <https://doi.org/10.1111/j.1365-3059.2009.02127.x>
- NSW (New South Wales) Government. 2012. Plant Biosecurity Orange. Exotic Pest Alert: *Potato Spindle Tuber Viroid* (PSTVd) in tomatoes. Department of primary Industries. Plant Biosecurity Primefact 1151: 4.
- Olmedo-Velarde A, Avijit R, Belanger CA, Watanabe S, Hamasaki RT, Mavrodieva V A, Nakhla MK and Melzer MJ. 2018. First report of Tomato chlorotic dwarf viroid infecting greenhouse tomato in Hawaii. *Plant Disease* “First Look” paper. <http://dx.doi.org/10.1094/PDIS-08-18-1401-PDN>
- Owens RA. 2007. *Potato spindle tuber viroid*: The simplicity paradox resolved? *Molecular Plant Pathology*, 8:549-560.



- Pérez-Moreno L, Rico-Jaramillo E, Sánchez-Palé JR, Ascencio-Ibáñez JT, Díaz-Plaza R y Rivera-Bustamante RF. 2004. Identificación de virus fitopatógenos en cultivos hortícolas de importancia económica en el estado de Guanajuato, México. *Revista Mexicana de Fitopatología* 22:187-197. <https://www.redalyc.org/pdf/612/61222205.pdf>
- Pieterse CMJ, Zamioudis C, Berendsen RL, Weller DM, Van Wees SCM and Bakker PAHM. 2014. Induced Systemic Resistance by Beneficial Microbes. *Annual Review of Phytopathology* 52:347–75.
- Puchta H, Herold T, Verhoeven K, Roenhorst A, Ramm K, Schmidt-Puchta W and Sanger, HL. 1990. A new strain of *potato spindle tuber viroid* (PSTVd-N) exhibits major sequence differences as compared to all other PSTVd strains sequenced so far. *Plant Molecular Biology* 15: 509–511.
- Ramos-Villanueva CD, Carrillo-Benitez G, Zamora-Macorra EJ, Santiago-Elena E, Ramrez-Alarcn S, Jimenez-Vidals J y vila Lpez R. 2023. Uso de microorganismos endfitos para el manejo del *Tomato brown rugose fruit virus* en cultivo de jitomate (*Solanum lycopersicum*). *Revista Mexicana de Fitopatologa*. 41(4): 109-117. <https://doi.org/10.18781/R.MEX.FIT.2023-1>
- Ramrez-Rojas S. 2006. Modelos para predecir rendimientos y prdidas debido a dao de virus en jitomate (*Lycopersicon esculentum* Mill). Tesis de doctorado. Colegio de Postgraduados. Montecillo, Texcoco, Edo. De Mxico. 50 p.
- Raupach GS, Liu L, Murphy JF, Tuzun S and Kloepper JW. 1996. Induced systemic resistance in cucumber and tomato against *Cucumber mosaic cucumovirus* using plant growth-promoting rhizobacteria (PGPR). *Plant Disease* 80:891–894. <https://doi.org/10.1094/pd-80-0891>
- Rojas A, Kvarnheden A, Marcenaro D and Valkonen JPT. 2005. Sequence characterization of Tomato leaf curl Sinaloa virus and Tomato severe leaf curl virus: Phylogeny of New World begomoviruses and detection of recombination. *Archives of Virology* 150: 1281-1299. [10.1007/s00705-005-0509-x](https://doi.org/10.1007/s00705-005-0509-x)
- Rojas MR, Acedo MA, Maliano MR, Soto-Aguilar M, Souza JO, Briddon RW, Kenyon L, Rivera-Bustamante RF, Murilo ZF, Adkins S, Legg JP, Kvarnheden A, Wintermantel WM, Sudarshana MR, Peterschmitt M, Lapidot M, Martin DP, Moriones E, Inoue-Nagata AK and Gilbertson RL. 2018. World management of Geminiviruses. *Annual Review of Phytopathology* 65:637-677. <https://doi.org/10.1146/annurev-phyto080615-10032>
- Ryals J, Uknes S and Ward E. 1994. Systemic acquired resistance. *Plant Physiology*. 104:1109–Rivarez MPS, Vucurovic A, Mehle N, Ravnikar M and Kutnjak D. 2021. Global Advances in Tomato Virome Research: Current Status and the Impact of High-Throughput Sequencing. *Frontier Microbiology* 12:671925. <https://doi.org/10.3389/fmicb.2021.671925>
- Sabanadzovic S, Valverde RA, Brown JK, Manrtin RR and Tzanetakis IE. 2009. Southern tomato virus: The link between the families Totiviridae and Partitiviridae. *Virus Research* 140(1-2): 130-137. <https://doi.org/10.1016/j.virusres.2008.11.018>
- Samaniego GB, Reyes RA, Moreno VOA y Tun SJM. 2017. Resistencia sistmica inducida contra virus fitoptgenos mediada por la inoculacin con la rizobacteria *Bacillus* spp. *La Revista de Proteccin Vegetal* 32(1):10-22.
- Snchez-del Castillo, F.E., Moreno-Prez C., Cruz-Arellanes, E.L. 2009. Produccin de jitomate hidropnico bajo invernadero en un sistema de dosel en forma de escalera. *Revista Chapingo Serie Horticultura*, 15:67-73.
- Sastry KS. 2013a. Seed-borne Plant Virus Diseases. Springer. India. 317 p. <https://doi.org/10.1007/978-81-322-0813-6>
- Sastry KS. 2013b. Plant Virus and Viroid Diseases in the Tropics. Introduction of Plant Viruses and Sub-Viral Agents, Classification, Assessment of Loss, Transmission and Diagnosis. Vol. I. Springer <https://doi.org/10.1007/978-94-007-6524-5>
- Savary S, Willocquet L, Pethybridge S, Esker J, McRoberts P and Nelson A. 2019. The global burden of pathogens and pests on major food crops. *Nature Ecology & evolution* 3(3): 430-439. <https://doi.org/10.1038/s41559-018-0793-y>
- Singh RP and Teixeira da Silva JA. 2006. Ornamental plants: Silent carrier of evolving viroids. In: J. A. Teixeira da Silva (ed.), *Floriculture, ornamental and plant biotechnology*, 3: 531–539.
- Singh RP, Xianzhou N and Singh M. 1999. Tomato chlorotic dwarf viroid: An evolutionary link in the origin of pospiviroids. *Journal of General Virology*, 80: 2823–2828.
- Singh RP, Ready KFM and Nie X. 2003. Part II. Biology. In ‘Viroids’. (Eds A Hadidi, R Flores, JW Randies, JS Semancik) pp. 30–48. (CSIRO Publishing: Australia).
- Singh RP and Dilworth DA. 2009. *Tomato chlorotic dwarf viroid* in the ornamental plant *Vinca minor* and its transmission through tomato seed. *European Journal of Plant Pathology*, 123(1), 111–116.

- Sofy AR, Sofy MR, Hmed AA and El-DougDoug K. 2019. Potential Effect of Plant Growth-Promoting Rhizobacteria (PGPR) on Enhancing Protection Against Viral Diseases. In: Maheshwari, D. K. and Dheeman, S. (eds). *Field Crops: Sustainable Management by PGPR*. Springer. pp 411-445.
- Suganyadevi M, Manoranjitham SK, Senthil N, Raveendran M and Karthikeyan G. 2018. Prevalence of Bud Blight of Tomato Caused by *Groundnut bud necrosis virus* in Tamil Nadu, India. *International Journal of Current Microbiology and Applied Sciences* 7(11): 734-742. e <https://doi.org/10.20546/ijcmas.2018.711.088>
- Sharaf ASAS, Taheri H, Ghodoum PMH A and Soleymani F. 2023. Molecular and phenotypic Responses of Rhizobacteria-Treated tomato plants to *Tomato mosaic virus* under greenhouse conditions. *Iran Journal Biotechnology* 21(1): e3220. <https://doi.org/10.30498/ijb.2022.319382.3220>
- Shirasu K, Nakajima H, Rajasekhar VK, Dixon RA and Lamb C. 1997. Salicylic acid potentiates an agonist-dependent gain control that amplifies pathogen signals in the activation of defense mechanisms. *Plant Cell* 9:261-270. <https://doi.org/10.1105/tpc.9.2.261>
- Scholthof KBG, Adkins S, Czosnek H, Palukaitis P, Jacquot E, Honh T, Hohn B, Saunders K, Candresse T, Ahlquist P, Hemenway C and Foster GD. 2011. Top 10 plant viruses in molecular plant pathology. *Molecular Plant Pathology* 12(9): 938-954. <https://doi.org/10.1111/j.1364-3703.2011.00752.x>
- Steyer S, Olivier T, Skelton A, Nixon T and Hobden E. 2010. *Columnnea latent viroid* (CLVD): first report in tomato in France. *Plant pathology* 59 (4): 794. <https://doi.org/10.1111/j.1365-3059.2010.02261.x>
- Taniguchi M, Sekine KT and Koeda. 2023. Lisianthus enation leaf curl virus, a begomovirus new to Japan, is more virulent than the prevalent tomato yellow leaf curl virus in Ty-gene-mediated resistant tomato cultivars. *Journal of General Plant Pathology* 89:35-46. <https://doi.org/10.1007/s10327-022-01101-5>
- van Brunschot SL, Verhoeven JThJ, Persley DM, Geering ADW, Drenth A and Thomas JE. 2014. An outbreak of Potato spindle tuber viroid in tomato is linked to imported seed Eur J Plant Pathol. <https://doi.org/10.1007/s10658-014-0379-8>
- Vanthana M, Nakkeeran S, Malathi VG, Renukadevi P and Vinodkumar S. 2019. Induction of in planta resistance by flagellin (*Flg*) and elongation factor-TU (*EF-Tu*) of *Bacillus amyloliquefaciens* (VB7) against groundnut bud necrosis virus in tomato. *Microbial Pathogenesis* 137: 103757
- Verhoeven JJJ, Jansen CCC, Willemsen TM, Kox LFF, Owens RA and Roenhorst JW. 2004. Natural infections of tomato by *Citrus exocortis viroid*, *Columnnea latent viroid*, *Potato spindle tuber viroid* and *Tomato chlorotic dwarf viroid*. *European Journal of Plant Pathology*, 110: 823–831,
- Verhoeven JJJ, Jansen CCC, Roenhorst JW, Flores R and de la Peña M. 2009. *Pepper chat fruit viroid*: Biological and molecular properties of a proposed new species of the genus Pospiviroid. *Virus Research* 144(1–2): 209-214. <https://doi.org/10.1016/j.virusres.2009.05.002>
- Verhoeven JJJ, Botermans M, Meekes ET, Meekes M and Roenhorst JW. 2012. *Tomato apical stunt viroid* in the Netherlands: most prevalent pospiviroid in ornamentals and first outbreak in tomatoes. *Eur J Plant Pathol* 133, 803–810 <https://doi.org/10.1007/s10658-012-0005-6>
- Verhoeven J Th J, J W Roenhorst, Owens, R. A. 2011. Mexican papita viroid and tomato planta macho viroid belong to a single species in the genus Pospiviroid. *Archives of Virology* 156(8):1433-7. [10.1007/s00705-011-0975-2](https://doi.org/10.1007/s00705-011-0975-2)
- Verhoeven JT, Jansen JCC, Willemsen TM, Kox LFF, Owens RA and Roenhorst JW. 2004. Natural infections of tomato by *Citrus exocortis viroid*, *Columnnea latent viroid*, *Potato spindle tuber viroid* and *Tomato chlorotic dwarf viroid*. *European Journal of Plant Pathology* 110: 823-831
- Verbeek M, Dulleman AM, van den Heuvel JFJM, Marris PC and van der Vlugt RAA. 2008. *Tomato marchitez virus*, a new plant picorna-like virus from tomato related to tomato torrado virus. *Archives of Virology* 153(1): 127-134. [10.1007/s00705-007-1076-0](https://doi.org/10.1007/s00705-007-1076-0)
- Vessey J K. 2003. Plant growth promoting rhizobacteria as biofertilizers. *Plant and Soil*. 255:571–586. <https://doi.org/10.1023/a:1026037216893>
- Walter B. 1987. Tomato apical stunt. pp. 321-327. In: Diener, T.O. (ed.). *The Viroids*. Plenum Press, New York, USA.
- Walter B, Thouvenal JC and Fauquet C. 1980. Les virus de la tomate en Cote d'Ivoire. *Annales de Phytopathologie* 12: 259– 275.

- Walters D. 2009. Are plants in the field already induced? Implications for practical disease control. *Crop Protection*. 28: 459-465.
- Wang S, Wu H, Qiao J, Ma L, Liu J, Xia Y and Gao X. 2009. Molecular mechanism of plant growth promotion and induced systemic resistance to *Tobacco mosaic virus* by *Bacillus* spp. *Journal of Microbiology and Biotechnology*. 19:1250–1258. <https://doi.org/10.4014/jmb.0901.008>
- Yanagisawa H and Matsushita Y. 2017. Host ranges and seed transmission of *Tomato planta macho viroid* and *Pepper chat fruit viroid*. *European Journal of Plant Pathology* 149:211–217. <https://doi.org/10.1007/s10658-017-1160-6>.
- Yanagisawa H and Matsushita Y. 2018. Differences in dynamics of horizontal transmission of Tomato planta macho viroid and Potato spindle tuber viroid after pollination with viroid-infected pollen. *Virology* 516: 258-264.
- Zehnder GW, Murphy JF, Sikora EJ, Kloepper JW. 2001. Application of rhizobacteria for induced resistance. *European Journal of Plant Pathology* 107:39–50. <https://doi.org/10.1023/a:1008732400383>
- Zehnder GW, Yao C, Murphy JF, Sikora ER and Kloepper JW. 2000. Induction of resistance in tomato against *Cucumber mosaic cucumovirus* by plant growth-promoting rhizobacteria. *Biocontrol* 45:127–137. <https://doi.org/10.1023/a:1009923702103>
- Zuñiga-Romano MC, Ochoa-Martínez DL, Rojas-Martínez RI and Zamora-Macorra EJ. 2019. Identification of begomoviruses in tomato and determination of seed transmission in Atlacomulco, Mexico State. *Mexican Journal of Phytopathology* 37(2): 279-286. <https://doi.org/10.18781/R.MEX.FIT.1902-2>
- Zhu L, Li Y, Li L, Yang J and Zhang M. 2011. Ethylene is involved in leafy mustard systemic resistance to Turnip mosaic virus infection through the mitochondrial alternative oxidase pathway. *Physiological and Molecular Plant Pathology* 76(3-4):166-172